Table 1. Performance of backgrounding calves fed a ration either as a total mixed ration (TMR; feed ingredients were mixed and delivered daily in a feed bunk) or as an ensiled ration (ingredients were mixed, ensiled, and offered in a self-feeding system; SF).

| | Treatments | | | |
|---|------------|-------|------------------|-----------------|
| Item | TMR | SF | SED ¹ | <i>P</i> -value |
| Initial BW, kg | 224 | 223 | 1.6 | 0.37 |
| Final BW, kg | 299 | 308 | 2.9 | 0.03 |
| DMI, kg/d | 7.37 | 8.59 | 0.193 | < 0.01 |
| DMI:BW, % | 2.91 | 3.24 | 0.062 | < 0.01 |
| ADG, kg/d | 1.07 | 1.22 | 0.039 | < 0.01 |
| ADG:DMI | 0.139 | 0.147 | 0.0029 | 0.36 |
| Coefficient of variation, % | | | | |
| Final BW | 5.3 | 5.3 | 0.68 | 0.95 |
| ADG | 14.5 | 14.7 | 2.18 | 0.93 |
| Δ BW coefficient of variation ² , percent units | 1.20 | 1.19 | 0.706 | 0.98 |

¹Standard error of the difference between means

²Final BW coefficient of variation minus Initial BW coefficient of variation