

Parameter	LEN + PEMBRO (N=355)	SUN (N=357)
Patients receiving any subsequent systemic anticancer therapy ^a , n (%)	117 (33.0)	206 (57.7)
Anti-VEGF	108 (30.4)	120 (33.6)
PD-1/PD-L1 checkpoint inhibitor	29 (8.2)	154 (43.1)
MTOR inhibitor	6 (1.7)	17 (4.8)
CTLA-4 inhibitor	6 (1.7)	18 (5.0)
Other	12 (3.4)	20 (5.6)
Median (range) time to next line therapy ^b , mos	12.68 (1.45–37.36)	6.62 (0.39–28.52)
Median (range) duration of first subsequent anticancer therapy ^c , mos	5.16 (0.10–30.23)	6.82 (0.03–30.72)
PFS2, median (95% CI)	Not reached (NE–NE)	28.7 mos (23.0–NE)
PFS2 HR (95% CI)	0.50 (0.39–0.65)	
Nominal <i>P</i> value	<0.0001	
PFS2 rate at 24/36 mos, % (95% CI)	72.7 (67.3, 77.4) / 61.9 (53.7, 69.0)	54.2 (48.4, 59.6) / 42.9 (32.8, 52.5)
^a Monotherapy or in combination; ^b includes patients with available start date of first subsequent systemic anticancer medication; ^c includes patients with available start and end dates of first subsequent systemic anticancer medication. CI, confidence interval; HR, hazard ratio; LEN, lenvatinib; mos, months; NE, not estimable; PEMBRO, pembrolizumab; PFS2, progression-free survival on next-line therapy; SUN, sunitinib.		