

TABLE 1. Patient and tumor characteristics for paratesticular liposarcoma (n = 60)

Characteristic	n	%/range
Median age at presentation (years)	66.5	
Prior cancer history		
Yes	12	20
No	48	80
Tumor anatomic location		
Paratesticular	42	70
Inguinal canal	10	17
In spermatic cord	8	13
Tumor laterality		
Right	31	52
Left	29	48
Initial surgery		
Yes	32	53
No	28	47
Diagnosis on initial presentation		
Concern for malignancy	28	47
Inguinal hernia	16	27
Unknown mass	11	18
Scrotal swelling	4	7
Hydrocele	1	2
Median tumor size (cm)	9.1	1.4 - 27
Pathology		
Well differentiated	22	37
Dedifferentiated	36	60
Pleomorphic	1	2
Myxoid	1	2
Surgery margin status after primary resection (n=55)		
Negative microscopic margins	20	36
Positive microscopic margins	31	56
Unknown	4	7
Radiation recommended		
Yes	19	32
No	41	68
Radiation timing to primary resection (n=19)		
Preoperative	6	32
Postoperative	10	53
Recurrent disease after primary resection		
Local	21	35
Distant	4	7
None	35	58
Median time to recurrence (months)	23.9	2.4 - 177.5
Median follow up (months)	48.5	1.5 - 209.5
Status at last follow up		
Alive with disease	11	18
Alive without disease	42	70
Deceased secondary to disease	5	8
Deceased for other cause	2	3